

PORIFERA

- Level of body organization ?
- Middle layer = ?
Acellular matrix = location of structural elements (spicules & spongin) & has cells moving through it = archeocytes
- Diagnostic cell type: ?

PORIFERA

- **CELLULAR** level of body organization
- Middle layer = **MESOHYL**
Acellular matrix = location of structural elements (spicules & spongin) & has cells moving through it = archeocytes
- Diagnostic cell type: **CHOANOCYTE**
= flagellated collar cell

The Three TYPES of Sponges



**Asconoid =
smallest**




**Syconoid =
middle-sized**

**Leuconoid
= Largest**



In the jar, these sponge specimens look like white or transparent plant roots..



PHYLUM Porifera
TYPE ?

In lab you could only look at a whole specimen (as above) in a jar or at prepared slides.

PHYLUM Porifera
TYPE Asconoid

NOTE: Many of our slide specimens have been stained red or green.
(Look like.....???????)

This is the smallest and simplest sponge type.
(i.e. they are too small to dissect.) Name often used for this most primitive unit?



PHYLUM Porifera
TYPE Asconoid



BSU - Basic Sponge Unit.

Choanocytes are located in the spongocoel.

What is the function of a gemmule?



PHYLUM Porifera

TYPE ?




Name this hole?

What is this?



PHYLUM Porifera
TYPE Asconoid



Note: Spicules at the neck of the osculum

Bud →

Terms you need to know: **spicules**, **spongocoel**, **bud** & **Osculum**. Compare to fig 1.3-A in your lab manuals.


Incurrent Pores (Ostia), Porocytes and Prosopyles

- Incurrent pores or ostia are the openings through which water first enters a sponge. These are usually formed by several cells.
- The PROSOPYLE is the name given to the entry hole/channel/pore leading into the area of choanocytes. It is formed by one donut-shaped cell, the porocyte.

Asconoid Sponges

As an incurrent pore or ostium, this opening brings water directly into the sponge.

It also serves as a **prosopyle**, bringing **water into contact with the choanocytes** lining the **spongocoel**.




Thus it has a dual function.

Thus the incurrent pore or ostium is serves as a **prosopyle**.

The actual opening is formed by 1 cell, the porocyte.

Syconoid Sponges

The ostia/incurrent pores in syconoid sponges are generally made of several cells. Water enters the sponge through these pores and moves into the incurrent canal.




Note the prominent spicules




PHYLUM Porifera
TYPE Syconoid

The choanocytes are located in the radial canals.
These are the 'middle-sized' sponges.



Water flow: Ostium -> Incurrent canal (I) -> Prosopyle channel (P) (a porocyte) -> Radial canals (R) (area of choanocytes) -> Apopyle channel (A) -> Spongocoel (S) -> Osculum (O)



Choanocytes
are
located
where?




PHYLUM Porifera
TYPE Leuconoid

The choanocytes are located in the many flagellated chambers.

Leuconoid Sponges

The ostia (several cells) allow water to enter incurrent canals.

Water leaves these to enter the flagellated chambers (**area of choanocytes**) via the prosopyles (**porocytes**)



Sponge Reproduction

Sponges are usually monoecious but can be dioecious

ASEXUAL

Marine

- Budding
- Fragmentation
- Regeneration

Freshwater sponges

- Gemmules
- + 3 methods above

SEXUAL

- Male & female gametes are formed.

Archeocytes become eggs

Choanocytes filter sperm out of the water

- Fertilization is involved.
- Planktonic larvae or mini flagellated colonies are released to colonize new areas.